The Cox model: introduction and history

Bianca L. De Stavola Centre for Statistical Methodology LSHTM

March 8, 2013





Today is a celebration of an incredibly influential paper:

- the most cited paper in the whole history of JRSS
- the third most cited paper in medical journals
- it has a total of nearly 30,000 citations (according to Web of Science)

References

and this is still increasing





・ロン ・四 と ・ ヨン ・ ヨン

Introduction	In 1972	Paper	Thanks	References	
Outline					



- 1 Introduction
- 2 In 1972 ...
- 3 'Regression Models and Life-Tables'
 - The Cox model
 - Insights
 - What was new
- 4 Trail of influence
 - The Discussion
 - The next 10 years
 - The next 20 years
 - Applications

5 Thanks

5/45

The next 10 years The next 20 years

Applications

4 Trail of influence The Discussion

- Insights
- What was new
- 3 'Regression Models and Life-Tables' The Cox model
- 1 Introduction
- 2 In 1972 ...





Thanks



In 1972



We are concerned with studying:

- individuals at risk of experiencing a failure after time T
- measured from a relevant origin and according to a relevant measurement scale
- difficulty if some are not observed until failure occurs, *i.e.* are censored
- crucially, censoring must be independent of the failure process



If T is a continuous positive random variable its probability distribution is equivalently specified by:

• the density function:

$$f(t) = \lim_{\Delta t \to 0^+} rac{\Pr\left(t \le T < t + \Delta t
ight)}{\Delta t}$$

• the survivor function:

$$S(t) = Pr(T \ge t)$$

the hazard function:

$$\lambda(t) = \lim_{\Delta t \to 0^+} rac{\Pr\left(t \le T < t + \Delta t | t \le T
ight)}{\Delta t}$$





By the product law of probability, S(t) is related to $\lambda(t)$:

$$S(t) = \lim \prod_{k=0}^{r-1} \{1 - \lambda(\tau_k)(\tau_{k+1} - \tau_k)\}$$

where:

- the limit is for $(\tau_{k+1} \tau_k) \rightarrow 0$
- $0 < \tau_1 < \tau_2 < \ldots < \tau_r = t$ the interval endpoints

▲□▶ ▲□▶ ▲臣▶ ★臣▶ 臣 のへで





By the product law of probability, S(t) is related to $\lambda(t)$:

$$S(t) = \lim \prod_{k=0}^{r-1} \left\{ 1 - \lambda(\tau_k) (\tau_{k+1} - \tau_k) \right\}$$

where:

• the limit is for
$$(\tau_{k+1} - \tau_k) \rightarrow 0$$

• $0 < \tau_1 < \tau_2 < \ldots < \tau_r = t$ the interval endpoints

S(t) is the product of the conditional survival probabilities for infinitesimal intervals up to t





By the product law of probability, S(t) is related to $\lambda(t)$:

$$S(t) = \lim \prod_{k=0}^{r-1} \left\{ 1 - \lambda(\tau_k) (\tau_{k+1} - \tau_k) \right\}$$

where:

• the limit is for
$$(\tau_{k+1} - \tau_k) \rightarrow 0$$

• $0 < \tau_1 < \tau_2 < \ldots < \tau_r = t$ the interval endpoints

S(t) is the product of the conditional survival probabilities for infinitesimal intervals up to t

construction of the likelihood depends on this

Three main analytical approaches:

1 Non-parametric estimation of S(t)

Trail

Thanks

References

- **2** Parametric estimation of S(t)
- 3 Comparison of survivor functions ("the two-sample problem")





Actuarial (Life-Table) estimation:

- long tradition in demography
- assuming hazard function piecewise constant over pre-specified intervals {t_j, t_{j+1}}, Â_j = no. events / total follow-up time
- survivor function estimated as the product of the conditional probabilities of surviving each interval: $\hat{S}(t_j) = \prod_{k < j} (1 \hat{\lambda}_k)$

イロン イロン イヨン イヨン 三日





Actuarial (Life-Table) estimation:

- long tradition in demography
- assuming hazard function piecewise constant over pre-specified intervals {t_j, t_{j+1}}, Â_j = no. events / total follow-up time
- survivor function estimated as the product of the conditional probabilities of surviving each interval: $\hat{S}(t_j) = \prod_{k < j} (1 \hat{\lambda}_k)$
- Product Limit estimation:
 - exactly the same but defined for vanishingly small intervals
 - hence $\hat{\lambda}_j =$ no. events / total no. persons at risk
 - derived by Kaplan & Meier (1958) as a non-parametric MLE of S(t)
- Although both derived from ML arguments, asym. properties not developed until later (Breslow and Crowley, 1974)

・ロト ・四ト ・ヨト ・ヨト 三田



- **Exponential and Weibull often used** (simple formulæ for S(t) and $\lambda(t)$)
- Approach attractive because of physical interpretation, e.g.

- multi-hit carcinogenesis theories lead to Weibull models (Armitage and Doll, 1954, 1961)
- Estimation via ML mostly derived assuming fixed censoring time (*e.g.* Bartholomew 1963 for exponential, Pike 1966 for Weibull)
- Inclusion of explanatory variables rare and with no censoring



Generalizations of the Savage–Wilcoxon rank test to settings with censored data:

- Mantel (1966):
 - test based on difference between observed and expected events at each failure time
 - expectations come from the hypergeometric distribution
 - results combined as in the Mantel-Haenszel test (1959)



Generalizations of the Savage–Wilcoxon rank test to settings with censored data:

- Mantel (1966):
 - test based on difference between observed and expected events at each failure time
 - expectations come from the hypergeometric distribution
 - results combined as in the Mantel-Haenszel test (1959)
- Peto and Peto (1972):
 - different derivation of the same comparison
 - named log-rank test
 - Note: paper read < 2 months before "Regression models and life-tables"

Introduction In 1972 ... Paper Trail Thanks References Summary of methods prevalent in 1972



- Statistical theory for non-parametric estimation of S(t) not yet fully formalized
- Inference for parametric estimation of S (t): complex even with simple censoring mechanisms
- Comparison of survivor curves dealt with via significance tests (and only possible for categorical variables)
- Extension of parametric models to include explanatory variables not generally available with censoring

Thanks



1 Introduction

- 2 In 1972 ...
- 3 'Regression Models and Life-Tables'
 - The Cox model
 - Insights
 - What was new
- 4 Trail of influence
 The Discussion
 The next 10 years
 The next 20 years
 Applications

5 Thanks

Regression Models and Life-Tables

References

By D. R. Cox

Imperial College, London

[Read before the ROYAL STATISTICAL SOCIETY, at a meeting organized by the Research Section, on Wednesday, March 8th, 1972, Mr M. J. R. HEALY in the Chair]

SUMMARY

The analysis of censored failure times is considered. It is assumed that on each individual are available values of one or more explanatory variables. The hazard function (age-specific failure rate) is taken to be a function of the explanatory variables and unknown regression coefficients multiplied by an arbitrary and unknown function of time. A conditional likelihood is obtained, leading to inferences about the unknown regression coefficients. Some generalizations are outlined.

Read here in the Goldsmiths Lecture Theatre

イロン イロン イヨン イヨン 三日

Trail



"The present paper is largely concerned with the extension of the results of Kaplan and Meier to the comparison of life tables..."

Thanks

- "... and more generally to the incorporation of regression like arguments into life-table analysis"
- it would be "sensible to make a minimum of assumptions leading to a convenient analysis"



- "The present paper is largely concerned with the extension of the results of Kaplan and Meier to the comparison of life tables..."
- "... and more generally to the incorporation of regression like arguments into life-table analysis"
- it would be "sensible to make a minimum of assumptions leading to a convenient analysis"



- "The present paper is largely concerned with the extension of the results of Kaplan and Meier to the comparison of life tables..."
- "... and more generally to the incorporation of regression like arguments into life-table analysis"
- it would be "sensible to make a minimum of assumptions leading to a convenient analysis"

イロン イロン イヨン イヨン 三日

Thanks



1 Introduction

- 2 In 1972 ...
- 3 'Regression Models and Life-Tables'The Cox model
 - Insights
 - What was new

4 Trail of influence■ The Discussion

- The next 10 years
- The next 20 years
- Applications

5 Thanks





- Let $\mathbf{z} = z_1, z_2, \dots, z_p$ be explanatory variables of interest
- Proportional hazards (PH) model defined as

$$\lambda(t; \mathbf{z}) = \exp(\mathbf{z}\boldsymbol{\beta}) \lambda_0(t)$$

β vector of unknown parameters (of interest)
 λ₀(t) unknown arbitrary function (nuisance)
 λ₀(t) describes the shape of the survival function
 exp(zβ) could be replaced by h(z, β)





- Let $\mathbf{z} = z_1, z_2, \dots, z_p$ be explanatory variables of interest
- Proportional hazards (PH) model defined as

$$\lambda(t; \mathbf{z}) = \exp(\mathbf{z}\boldsymbol{\beta}) \lambda_0(t)$$

β vector of unknown parameters (of interest)
 λ₀(t) unknown arbitrary function (nuisance)
 λ₀(t) describes the shape of the survival function
 exp(zβ) could be replaced by h(z, β)

- explore the consequences of allowing $\lambda_0(t)$ to be arbitrary
- method to have sensible properties, whatever $\lambda_0(t)$
- plausible loss of information about β is usually slight





Observations:

- n individuals, k fail
- independent censoring
- failure times: $0 < t_{(1)} < t_{(2)} < \ldots < t_{(k)} < \infty$
- $\Re(t)$ the set of individuals at risk at time t





- Observations:
 - n individuals, k fail
 - independent censoring
 - failure times: $0 < t_{(1)} < t_{(2)} < \ldots < t_{(k)} < \infty$
 - $\Re(t)$ the set of individuals at risk at time t
- Originally estimation derived from a 'conditional likelihood'
 This is the product of factors, one per event time t_(i):

$$\frac{\exp \mathbf{z}_{(i)}\boldsymbol{\beta}}{\boldsymbol{\Sigma}_{\ell\in\Re\left(\boldsymbol{t}_{(i)}\right)}\exp\left\{\mathbf{z}_{\ell}\boldsymbol{\beta}\right\}}$$





- Observations:
 - n individuals, k fail
 - independent censoring
 - failure times: $0 < t_{(1)} < t_{(2)} < \ldots < t_{(k)} < \infty$
 - $\Re(t)$ the set of individuals at risk at time t
- Originally estimation derived from a 'conditional likelihood'
 This is the product of factors, one per event time t_(i):

$$\frac{\exp \mathsf{z}_{(i)}\beta}{\Sigma_{\ell\in\Re\left(t_{(i)}\right)}\exp\left\{\mathsf{z}_{\ell}\beta\right\}}$$

conditional probabilities that the failure is on the individual as observed





$$I(\boldsymbol{\beta}) = \sum_{i=1}^{k} \mathbf{z}_{(i)} \boldsymbol{\beta} - \sum_{i=1}^{k} \log \left[\sum_{\ell \in \Re(t_{(i)})} \exp \left\{ \mathbf{z}_{\ell} \boldsymbol{\beta} \right\} \right]$$





$$I(\boldsymbol{\beta}) = \sum_{i=1}^{k} \mathbf{z}_{(i)} \boldsymbol{\beta} - \sum_{i=1}^{k} \log \left[\sum_{\ell \in \Re(t_{(i)})} \exp \left\{ \mathbf{z}_{\ell} \boldsymbol{\beta} \right\} \right]$$

• The score function:

$$U_{\xi}\left(\beta\right) = \sum_{i=1}^{k} \left\{ z_{\xi i} - A_{\xi i}\left(\beta\right) \right\}$$

where $A_{\xi i}\left(oldsymbol{eta}
ight)$ is the weighted average of z_{ξ} in \Re

イロト イロト イヨト イヨト 二日





$$I(\boldsymbol{\beta}) = \sum_{i=1}^{k} \mathbf{z}_{(i)} \boldsymbol{\beta} - \sum_{i=1}^{k} \log \left[\sum_{\ell \in \Re(t_{(i)})} \exp \left\{ \mathbf{z}_{\ell} \boldsymbol{\beta} \right\} \right]$$

• The score function:

$$U_{\xi}\left(\beta\right) = \sum_{i=1}^{k} \left\{ z_{\xi i} - A_{\xi i}\left(\beta\right) \right\}$$

where $A_{\xi i}\left(oldsymbol{eta}
ight)$ is the weighted average of z_{ξ} in \Re

The information matrix the sum of weighted covariance matrices





$$I(\boldsymbol{\beta}) = \sum_{i=1}^{k} \mathbf{z}_{(i)} \boldsymbol{\beta} - \sum_{i=1}^{k} \log \left[\sum_{\ell \in \Re(t_{(i)})} \exp \left\{ \mathbf{z}_{\ell} \boldsymbol{\beta} \right\} \right]$$

The score function:

$$U_{\xi}\left(\beta\right) = \sum_{i=1}^{k} \left\{ z_{\xi i} - A_{\xi i}\left(\beta\right) \right\}$$

where $A_{\xi i}\left(oldsymbol{eta}
ight)$ is the weighted average of z_{ξ} in \Re

- The information matrix the sum of weighted covariance matrices
- $\hat{\boldsymbol{\beta}}$ found by iteration





$$I(\boldsymbol{\beta}) = \sum_{i=1}^{k} \mathbf{z}_{(i)} \boldsymbol{\beta} - \sum_{i=1}^{k} \log \left[\sum_{\ell \in \Re(t_{(i)})} \exp \left\{ \mathbf{z}_{\ell} \boldsymbol{\beta} \right\} \right]$$

The score function:

$$U_{\xi}\left(\beta\right) = \sum_{i=1}^{k} \left\{ z_{\xi i} - A_{\xi i}\left(\beta\right) \right\}$$

where $A_{\xi i}\left(oldsymbol{eta}
ight)$ is the weighted average of z_{ξ} in \Re

- The information matrix the sum of weighted covariance matrices
- $\hat{\boldsymbol{\beta}}$ found by iteration

Derivation was controversial



• $\frac{\exp z_{(i)}\beta}{\sum_{\ell \in \Re(t_{(i)})} \exp\{z_{\ell}\beta\}}$: interpreted as cond. prob. that individual (*i*) is the one failing at time $t_{(i)}$, given that a failure occurs at $t_{(i)}$

Thanks

- but it is given $\Re(t_{(i)})$
- equivalent to conditioning on the history of the process up to t
- independent of times \Rightarrow conditional probabilities for the ranks: $l(\beta)$ is marginal log lik. of the ranks (Kalbfleisch and Prentice, 1973)
- Cox (1975) calls it Partial Likelihood (PL) and shows max PLE consistent and asym. normal, with asym. covariance matrix estimated consistently (ordering of t_(i) defines a nesting of conditioning events: U and C derived cond. but hold uncond.)
- Tsiatis (1981) shows this more formally using empirical processes



• $\frac{\exp z_{(i)}\beta}{\sum_{\ell \in \Re(t_{(i)})} \exp\{z_{\ell}\beta\}}$: interpreted as cond. prob. that individual (*i*) is the one failing at time $t_{(i)}$, given that a failure occurs at $t_{(i)}$

Thanks

- but it is given $\Re(t_{(i)})$
- equivalent to conditioning on the history of the process up to t
- independent of times \Rightarrow conditional probabilities for the ranks: $l(\beta)$ is marginal log lik. of the ranks (Kalbfleisch and Prentice, 1973)
- Cox (1975) calls it Partial Likelihood (PL) and shows max PLE consistent and asym. normal, with asym. covariance matrix estimated consistently (ordering of t_(i) defines a nesting of conditioning events: U and O derived cond. but hold uncond.)
- Tsiatis (1981) shows this more formally using empirical processes



• $\frac{\exp z_{(i)}\beta}{\sum_{\ell \in \Re(t_{(i)})} \exp\{z_{\ell}\beta\}}$: interpreted as cond. prob. that individual (*i*) is the one failing at time $t_{(i)}$, given that a failure occurs at $t_{(i)}$

Thanks

- but it is given $\Re(t_{(i)})$
- equivalent to conditioning on the history of the process up to t
- independent of times \Rightarrow conditional probabilities for the ranks: $I(\beta)$ is marginal log lik. of the ranks (Kalbfleisch and Prentice, 1973)
- Cox (1975) calls it Partial Likelihood (PL) and shows max PLE consistent and asym. normal, with asym. covariance matrix estimated consistently (ordering of t_(i) defines a nesting of conditioning events: U and 3 derived cond. but hold uncond.)
- Tsiatis (1981) shows this more formally using empirical processes



• $\frac{\exp z_{(i)}\beta}{\sum_{\ell \in \Re(t_{(i)})} \exp\{z_{\ell}\beta\}}$: interpreted as cond. prob. that individual (*i*) is the one failing at time $t_{(i)}$, given that a failure occurs at $t_{(i)}$

Thanks

- but it is given $\Re(t_{(i)})$
- equivalent to conditioning on the history of the process up to t
- independent of times \Rightarrow conditional probabilities for the ranks: $I(\beta)$ is marginal log lik. of the ranks (Kalbfleisch and Prentice, 1973)
- Cox (1975) calls it Partial Likelihood (PL) and shows max PLE consistent and asym. normal, with asym. covariance matrix estimated consistently (ordering of t_(i) defines a nesting of conditioning events: U and 3 derived cond. but hold uncond.)
- Tsiatis (1981) shows this more formally using empirical processes



 exp z_(i)β
 Σ_{ℓ∈ℜ(t_(i))} exp{z_ℓβ}: interpreted as cond. prob. that individual (i) is the one failing at time t_(i), given that a failure occurs at t_(i)

Thanks

References

- but it is given $\Re(t_{(i)})$
- equivalent to conditioning on the history of the process up to t
- independent of times \Rightarrow conditional probabilities for the ranks: $I(\beta)$ is marginal log lik. of the ranks (Kalbfleisch and Prentice, 1973)
- Cox (1975) calls it Partial Likelihood (PL) and shows max PLE consistent and asym. normal, with asym. covariance matrix estimated consistently (ordering of t_(i) defines a nesting of conditioning events: U and 3 derived cond. but hold uncond.)

Tsiatis (1981) shows this more formally using empirical processes

In 1972 Is it a conditional likelihood?

Paper

Trail



• $\frac{\exp z_{(i)}\beta}{\sum_{\ell \in \Re(t_{(i)})} \exp\{z_{\ell}\beta\}}$: interpreted as cond. prob. that individual (*i*) is the one failing at time $t_{(i)}$, given that a failure occurs at $t_{(i)}$

Thanks

References

- but it is given $\Re(t_{(i)})$
- equivalent to conditioning on the history of the process up to t
- independent of times \Rightarrow conditional probabilities for the ranks: $I(\beta)$ is marginal log lik. of the ranks (Kalbfleisch and Prentice, 1973)
- Cox (1975) calls it Partial Likelihood (PL) and shows max PLE consistent and asym. normal, with asym. covariance matrix estimated consistently (ordering of $t_{(i)}$ defines a nesting of conditioning events: U and \Im derived cond, but hold uncond.)
- Tsiatis (1981) shows this more formally using empirical processes

Thanks



1 Introduction

2 In 1972 ...

3 'Regression Models and Life-Tables'
 The Cox model

- Insights
- What was new

4 Trail of influence
The Discussion
The next 10 years
The next 20 years
Applications

Applications

5 Thanks



If there are tied events the PL is not appropriate. To deal with this the paper proposes two strategies:

Thanks

References

◆□▶ ◆□▶ ◆臣▶ ◆臣▶ = 臣 = のへ⊙

De Stavola/History · 8 March 2013





If there are tied events the PL is not appropriate. To deal with this the paper proposes two strategies:

- I If few, a correction of the term contributing to the partial likelihood
- If several, PH model replaced by a proportional odds (PO) model:

$$\frac{\lambda(t; \mathbf{z}) dt}{1 - \lambda(t; \mathbf{z}) dt} = \frac{\lambda_0(t) dt}{1 - \lambda_0(t) dt} \exp(\mathbf{z}\beta)$$

where $\lambda_{0}\left(t\right) = \Pr\left(T \leq t + 1 | T > t\right)$ arbitrary





If there are tied events the PL is not appropriate. To deal with this the paper proposes two strategies:

- If few, a correction of the term contributing to the partial likelihood
- If several, PH model replaced by a proportional odds (PO) model:

$$\frac{\lambda(t; \mathbf{z}) dt}{1 - \lambda(t; \mathbf{z}) dt} = \frac{\lambda_0(t) dt}{1 - \lambda_0(t) dt} \exp(\mathbf{z}\beta)$$

where $\lambda_{0}\left(t
ight)= extsf{Pr}\left(extsf{T}\leq t+1| extsf{T}>t
ight)$ arbitrary

Similar arguments lead to PLE PO model \Rightarrow PH model as the intervals become infinitesimal

Introduction In 1972... Paper Trail Thanks References Additional insights



Other insights, both methodological and relevant for applications: Dealing with the two sample problem:

- with PH model this becomes a comparison of $\lambda_0(t)$ and $e^{\beta_1}\lambda_0(t)$
- score test from the PL for discrete times equivalent to Mantel's test and asym. equivalent to log-rank test
- novelty: it can be applied to continuous exposures

Introduction In 1972 ... Paper Trail Thanks References Additional insights



Other insights, both methodological and relevant for applications: Dealing with the two sample problem:

- with PH model this becomes a comparison of $\lambda_0(t)$ and $e^{\beta_1}\lambda_0(t)$
- score test from the PL for discrete times equivalent to Mantel's test and asym. equivalent to log-rank test
- novelty: it can be applied to continuous exposures
- 2 Departures from proportionality:
 - formulation of both PH and PO allows for time varying explanatory variables
 - special case: covariate generated from the interaction between a time fixed variable and time
 - This allows testing the proportional assumption

(日) (四) (日) (日) (日) (日)

Introduction In 1972 ... Paper Trail Thanks References Additional insights



Other insights, both methodological and relevant for applications: Dealing with the two sample problem:

- with PH model this becomes a comparison of $\lambda_0(t)$ and $e^{\beta_1}\lambda_0(t)$
- score test from the PL for discrete times equivalent to Mantel's test and asym. equivalent to log-rank test
- novelty: it can be applied to continuous exposures
- 2 Departures from proportionality:
 - formulation of both PH and PO allows for time varying explanatory variables
 - special case: covariate generated from the interaction between a time fixed variable and time
 - This allows testing the proportional assumption

3 Estimating failure time dsn: generalization of the product limit estimation of $\lambda(t)$

Thanks



1 Introduction

2 In 1972 ...

3 'Regression Models and Life-Tables'
 The Cox model

- The Cox mode
- Insights
- What was new
- 4 Trail of influence
 The Discussion
 The next 10 years
 The next 20 years
 Applications





The paper is well known although maybe not so well read

- Aspects that are well known:
 - semi-parametric PH model and its estimation approach
 - ability to perform score tests for continuous exposures





The paper is well known although maybe not so well read

- Aspects that are well known:
 - semi-parametric PH model and its estimation approach
 - ability to perform score tests for continuous exposures
- Aspects that are not so well known:
 - checking of PH assumption
 - solutions for tied event times, including semi-parametric PO model for discrete times
 - estimation of cumulative failure time distribution
 - extension to multivariate T and links to the accelerated failure time models
 - physical versus empirical interpretation of the model

Applications

The next 10 years The next 20 years

- The Discussion
- 4 Trail of influence
- Insights What was new
- The Cox model

3 'Regression Models and Life-Tables'



Outline



Trail







Extremely influential paper methodologically on three 'time scales':

Thanks

References

- 1 at the time origin: the Society discussion
- 2 in the next 10 years
- 3 in the following 20 years and beyond

Thanks

References

1 Introduction

2 In 1972 ...

Outline

- 3 'Regression Models and Life-Tables'
 The Cox model
 Insights
 - Insights
 - What was new
- 4 Trail of influence
 The Discussion
 The next 10 years
 The next 20 years
 Applications

5 Thanks





Highlights of that discussion:

Richard Peto: proposed an alternative approach to dealing with tied events

Thanks

References

- 2 Jack Kalbfleish and Ross Prentice: raised questions regarding the 'conditional' likelihood
- 3 Norman Breslow: showed how the baseline cumulative hazard function could be estimated in a more natural way
- Susannah Howard: showed how easily max. PL estimation could be performed

Thanks

Trail

References

1 Introduction

Outline

- 3 'Regression Models and Life-Tables' The Cox model
 - Insights
 - What was new
- 4 Trail of influence The Discussion The next 10 years The next 20 years Applications



IntroductionIn 1972 ...Pape2 - The next 10 years



(a) The development of the theory of partial likelihood

Trail

Thanks

References

(b) The incorporation within counting processes theory:

- Andersen and Gill (1982) simplified and generalized the results on asym. properties of PLE using martingale theory for counting processes (from Aalen 1975)
- Indeed this viewpoint is required for an elegant derivation of these properties
- PH model played a key role for this powerful methodological development

・ロト ・四ト ・ヨト ・ヨト

IntroductionIn 1972 ...Paper2 - The next 10 years



(a) The development of the theory of partial likelihood

Trail

Thanks

References

- (b) The incorporation within counting processes theory:
 - Andersen and Gill (1982) simplified and generalized the results on asym. properties of PLE using martingale theory for counting processes (from Aalen 1975)
 - Indeed this viewpoint is required for an elegant derivation of these properties
 - PH model played a key role for this powerful methodological development

◆□ → < □ → < Ξ → < Ξ → Ξ の Q ↔ 33/45

The next 10 yearsThe next 20 years

Applications

5 Thanks

1 Introduction

- 2 In 1972 ...
- 3 'Regression Models and Life-Tables'
 The Cox model
 - Insights
 - What was new

4 Trail of influence

Outline



Trail

Thanks References



Introduction In 1972 ... Paper Trail Thanks 3 – The next 20 years and beyond



 Interest in semi-parametric models exploded following PH model, prompting huge developments in theory (Bickel et al, 1998)

References

 These developments are increasingly important in causal inference, missing data, etc. (Tsiatis, 2006)

Introduction In 1972 ... Paper Trail Thanks 3 – The next 20 years and beyond



 Interest in semi-parametric models exploded following PH model, prompting huge developments in theory (Bickel et al, 1998)

References

 These developments are increasingly important in causal inference, missing data, *etc.* (Tsiatis, 2006)

Would we be using doubly robust methods, efficient gestimation, targeted ML *etc.* had "Regression Models and Life-Tables" not been published?

Thanks

1 Introduction

Outline

- 3 'Regression Models and Life-Tables' The Cox model
 - Insights
 - What was new

4 Trail of influence

- The Discussion The next 10 years The next 20 years
- Applications



Trail

Trail

Thanks

References



- The paper states that the proposed methodology will be for "applications in industrial reliability studies and in medical statistics"
- Was this a fair prediction?

De Stavola/History · 8 March 2013

36/45

Introduction In 1972 ... Paper Trail Than Citations from Web of Science Area proportional to number of citations





References

◆□ > ◆□ > ◆臣 > ◆臣 > 善臣 - のへで

Introduction In 1972 Paper Trail Thank Citations from Web of Science Area proportional to number of citations



References



Introduction In 1972 ... Paper Trail Tha Citations from Web of Science Area proportional to number of citations



References



Sac 40/45

5 Thanks

The next 20 years Applications

The next 10 years

- 4 Trail of influence The Discussion
- What was new
- 3 'Regression Models and Life-Tables' The Cox model

Insights

- 1 Introduction



Outline







Votes of thanks aired on 8 March 1972:

As usual [David Cox's] statistical ideas are of both theoretical interest and great practical importance.

(F. Downton)







Votes of thanks aired on 8 March 1972:

As usual [David Cox's] statistical ideas are of both theoretical interest and great practical importance.

(F. Downton)

... he has opened up new territories to common sense. (R. Peto)

Introduction Source	n In 1972 2S	Paper Trail	Thanks	References	and the
	Andersen PK (Survival analys hazards regress <i>Statistics in M</i>	(1991) sis 1982–199 sion model. <i>ledicine</i> , 10:1	1: the se	econd decade of	the proportional
	Cox DR, Oake Analysis of Sur Chapman and	s D (1984) <i>rvival Data.</i> Hall: Londo	n.		

Kalbfleisch JD, Prentice RL (2000) The Statistical Analysis of Failure Time Data. John Wiley: Canada.

Oakes D (1991)

Life table analysis. In: *Statistical Theory and Modelling*, Hinkley DV, Reid N, Snell EJ (eds).

Chapman and Hall: London.





🕯 Aalen OO (1975)

Statistical inference for a family of counting processes. Ph.D. thesis, University of California, Berkeley.

Trail

Andersen PK, Gill RD (1982)
 Cox's regression model for counting processes: a large sample study.
 Annals of Statistics, 10:1100–1120.

Thanks

References

- Bickel et al (1998) Efficient and Adaptive Estimation for Semiparametric Models. Springer: New York.
- Breslow N, Crowley J (1974)

A Large Sample Study of the Life Table and Product Limit Estimates Under Random Censorship.

Annals of Statistics, 2:437-453.



JRSS B, 135:185-207.

Introduction In 1972 References (3)

Trail

References

Thanks



Tsiatis AA (1981)

A large sample study of Cox's regression model. Annals of Statistics, 9:93–108.

Tsiatis AA (2006)

Semiparametric Theory and Missing Data.

Springer: New York.